

# Impact® 250 SC

Fungicide

# **IMPACT® 250 SC FUNGICIDE AT A GLANCE**

Registration No. L9064

Active ingredients Flutriafol (Triazole) – 250 g/t

Chemical class Pyrimidine

FRAC Group 3

## **MODE OF ACTION**

Flutriafol is a broad-spectrum triazole fungicide used for the control of important cereal diseases including powdery mildews, rusts, *Septoria spp.*, and *Rhynchosporium secalis*.

- 1 It shares a common mode of action with other triazole fungicides by inhibiting the C-14 $\alpha$ -demethylase enzyme involved in the biosynthesis of fungal sterols.
- 2 A characteristic structural feature of flutriafol is the tertiary alcohol with a 1,2,4-triazolylmethyl group and two differently substituted fluorophenyl groups.

#### RESISTANCE MANAGEMENT

For resistance management, **Impact® 250 SC** is a group code 3 (triazole) fungicide. Any fungal pathogen population may contain individuals naturally resistant to **Impact® 250 SC** and other Group code 3 fungicides. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly and exclusively in programs. These resistant fungal pathogens may not be controlled by **Impact® 250 SC** or any other group code 3 fungicides. To delay fungicide resistance:

- Avoid exclusive repeated use of fungicides from the same group code. Alternate or tank mix with products from different fungicide group codes.
- Integrate other control methods (chemical, cultural, biological) into disease control programs.



USE PLANT PROTECTION PRODUCTS SAFELY AND WITH RESPONSIBLE CARE.
PLEASE ALWAYS FOLLOW THE LABEL WHEN APPLYING PLANT PROTECTION PRODUCTS.



## **CROPS AND PESTS CONTROLLED**

Refer to label for application rates and time of application.

CROPS	DISEASES CONTROLLED
Barley	Leafspot (Rhynchosporium secalis), Net (Leaf) blotch (Pyrenophora teres) (Helminthosporium) Powdery mildew (Erysiphe graminis), Leaf rust (Puccinia hordei)
Wheat	Eye spot (Pseudocercosporella herpotrichoides), Speckled leaf blotch (Septoria tritici), Glume blotch (Septoria nodorum), Powdery mildew (Erysiphe graminis), Leaf rust (Puccinia recondita), Yellow rust (Puccinia striiformis)
Maize & Sweet Corn	Northern leaf blight (Exserohilium turcicum)
Potatoes	Early blight (Alternaria solani)
Soybean	Rust (Phakopsora pachyrhizi)



## FOR MORE INFORMATION

Put this powerful tool to work for you. Contact your local crop protection retailer, or FMC representative to learn how you can get more consistent control of key diseases in various crops with **Impact® 250 SC** fungicide.

#### FMC Chemicals (Pty) Ltd

Company Registration Number: 1988/001451/07 West End Office Park, Building C, Cnr West Ave & Hall Street, Centurion, 0014 www.ag.fmc.com/za/en IMPACT® 250 SC Reg. No. L9064, Act No. 36 of 1947, contains flutriafol (Triazole) – 250 g/L, Warning, Causes serious eye irritation, Harmful if inhaled, May cause respiratory irritation, Toxic to aquatic life with long lasting effects.

FMC Chemicals (Pty) Ltd, PO Box 44, Postnet Menlyn, Waterkloof Glen, 0181, Republic of South Africa.

Impact is a  $\,$  trademark of FMC Corporation or its affiliates.